CONTENTS

	Page
A Teosinte-Maize Hybrid. G. N. COLLINS and J. H. KEMPTON	1
Banana Root-Borer. G. F. MOZNETTE	39
Effect of Calcium Sulphate on the Solubility of Soils. M. M. McCOOL and C. E. MILLAR	47
Further Studies on the Influence of Humidity upon the Strength and Elasticity of Wool Fiber. J. I. HARDY	55
Composition and Density of the Native Vegetation in the Vicinity of the Northern Great Plains Field Station. J. T. SARVIS	63
Effect of Reaction of Solution on Germination of Seed and on Growth of Seedlings. ROBERT M. SALTER and T. C. McILVAINE	73
Philippine Downy Mildew of Maize. WILLIAM H. WESTON, JR.	97
Effect of Drugs on Milk and Fat Production. FRANK A. HAYS and MERTON G. THOMAS	123
Artificial and Insect Transmission of Sugar-Cane Mosaic. E. W. BRANDES	131
Halo-Blight of Oats. CHARLOTTE ELLIOTT	139
Influence of Fermentation on the Starch Content of Experimental Silage. ARTHUR W. DOX and LESTER YODER	173
Effect of Premature Freezing on Composition of Wheat. M. J. BLISH	181
Behavior of the Citrus-Canker Organism in Soil. H. ATHERTON LEE	189
Decline of Pseudomonas citri in the Soil. H. R. FULTON	207
Variation of Individual Pigs in Economy of Gain. R. C. ASHBY and A. W. MALCOMSON	225
Production of Conidia in Gibberella saubinetii. JAMES G. DICKSON and HELEN JOHANN	235
Effect of Manure-Sulphur Composts upon the Availability of the Potassium of Greensand. A. G. McCALL and A. M. SMITH	239
Rust in Seed Wheat and Its Relation to Seedling Infection. CHARLES W. HUNGERFORD	257
Practical Universality of Field Heterogeneity as a Factor Influencing Plot Yields. J. ARTHUR HARRIS	279
Transmission of the Mosaic Disease of Irish Potatoes. E. S. SCHULTZ and DONALD FOLSOM	315
Relative Susceptibility to Citrus-Canker of Different Species and Hybrids of the Genus Citrus, Including the Wild Relatives. GEORGE L. PELTIER and WILLIAM J. FREDERICH	339

Page

Presoak Method of Seed Treatment: A Means of Preventing Seed Injury Due to Chemical Disinfectants and of Increasing Germicidal Efficiency. HARRY BRAUN 363
Daily Development of Kernels of Hannchen Barley from Flowering to Maturity at Aberdeen, Idaho. HARRY V. HARLAN 393
Development of Barley Kernels in Normal and Clipped Spikes and the Limitations of Awnless and Hooded Varieties. HARRY V. HARLAN and STEPHEN ANTHONY 431
Investigations in the Ripening and Storage of Bartlett Pears. J. R. MAGNESS 473
Further Data on the Orange Rusts of Rubus. L. O. KUNKEL 501
Germ-Free Filtrates as Antigens in the Complement-Fixation Test. WILLIAM S. GOCHENOUR 513
Mosaic Disease of Corn. E. W. BRANDES 517
Genetics of Rust Resistance in Crosses of Varieties of Triticum vulgare with Varieties of T. durum and T. dicoccum. H. K. HAYES, JOHN H. PARKER, and CARL KURTZWEIL 523
Line-Selection Work with Potatoes. O. B. WHIPPLE 543
Occurrence of the Fixed Intermediate, Hordeum intermedium haxtoni, in Crosses between H. vulgare pallidum and H. distichon palmella. HARRY V. HARLAN and H. K. HAYES 575
Index 593

ERRATA AND AUTHORS' EMENDATIONS

Page 97, line 20, "were" should read "would be."

Page 110, Table I, column 1, line 13, "24" should read "23."

Page 136, line 2, "*Pseudococcus boninensis*" should read "*Pseudococcus boninsis.*"

Page 202, following line 16, add "*C. trifoliata* seedlings, however, on testing are readily susceptible to the disease."

Page 232, Table XVI, footnote, "$r=-0.452\pm0.068$" should read "$r=-0.166\pm0.083$."

Page 236, line 1, "The spores, both conidia and ascospores, behaved alike in germination" should read "The conidia from both conidia and ascospores behaved alike in germination."

Page 242, line 22, "$Al_2(SO_4)_3$ 0.18 H_2O" should read "$Al_2(SO_4)_3$ .18 H_2O" and "$FeSO_4$ 0.7 H_2O" should read "$FeSO_4$ .7 H_2O." The same correction should be made in Tables III, IV, V, and VII.

Page 278, Plate 47 and legend. "Internal hilar sorus shown at *x*" should be omitted. The marking *x* is incorrectly placed on the plate. There are two internal hilar sori shown in the lower right-hand portion of the plate.

Page 362, legend for Plate 59, line 10, "Infected leaves from twigs" should read "Infected leaves and twigs."

ILLUSTRATIONS

A Teosinte-Maize Hybrid

Text Figures

Page

1. Height: frequency distribution of plants in F_2 13
2. Total leaves: frequency distribution of plants in F_2 13
3. Height of sucker: frequency distribution of plants in F_2 14
4. Sucker index: frequency distribution of plants in F_2 14
5. Circumference index: frequency distribution of plants in F_2 16
6. Nodes without branches: frequency distribution of plants in F_2 15
7. Nodes above: frequency distribution of plants in F_2 17
8. Nodes above on third: frequency distribution of plants in F_2 18
9. Nodes on third: frequency distribution of plants in F_2 18
10. Primary branches: frequency distribution of plants in F_2 19
11. Secondary branches: frequency distribution of plants in F_2 19
12. Secondary index: frequency distribution of plants in F_2 20
13. Tassel branches on third: frequency distribution of plants in F_2 20
14. Male branch index: frequency distribution of plants in F_2 21
15. Male secondaries: frequency distribution of plants in F_2 22
16. Double male alicoles: frequency distribution of plants in F_2 23
17. Mixed alicoles: frequency distribution of plants in F_2 23
18. Single female alicoles: frequency distribution of plants in F_2 24
19. Double female alicoles: frequency distribution of plants in F_2 24
20. Alicole index: frequency distribution of plants in F_2 25
21. Nodes silking on third: frequency distribution of plants in F_2 25
22. Nodes silking index: frequency distribution of plants in F_2 26
23. Prophyllary spikes: frequency distribution of plants in F_2 26
24. Length of prophyllary: frequency distribution of plants in F_2 27
25. Rows in central spike: frequency distribution of plants in F_2 27
26. Rows of alicoles: frequency distribution of plants in F_2 28
27. Position of best spike: frequency distribution of plants in F_2 29
28. Number of alicoles: frequency distribution of plants in F_2 29
29. Number of suckers: frequency distribution of plants in F_2 30
30. Branch silking first: frequency distribution of plants in F_2 31
31. Days to pollen: frequency distribution of plants in F_2 31
32. Pollen to silk: frequency distribution of plants in F_2 32
33. Length of internode on third: frequency distribution of plants in F_2 32

Plates

1. A.—General view of F_2 plants of teosinte-maize hybrid. B.—F_2 plants of teosinte-maize hybrid, showing diversity in size and season 38
2. Teosinte-maize hybrid: A.—F_2 plant No. 36. B.—F_2 plant No. 49 38
3. Teosinte-maize hybrid: Pistillate inflorescence of F_2 plant No. 36, shown in Plate 2, A 38
4. Pistillate inflorescence of plant No. 49, shown in Plate 2, B 38
5. Pistillate inflorescences from plant No. 94, illustrating an intermediate type of inflorescence 38

Page

6. Teosinte-maize hybrid: A.—F_2 plant No. 31, showing compact growth characteristic of many of the plants. B.—F_2 plant No. 113, showing stiff, erect leaves. C.—F_1 plant, grown at Lanham, Md. 38
7. Teosinte-maize hybrid: Pistillate inflorescences of the F_1 plant shown in plant 6, C 38

BANANA ROOT-BORER

Text Figure

1. *Cosmopolites sordidus:* Section of sheath with egg *in situ* in compartment. 44

Plates

8. Banana root-borer (*Cosmopolites sordidus*): Adult 46
9. Egg and larva of banana root-borer: A.—Egg. B.—Larva, side view. C.—Head of larva, dorsal view. D.—Head of larva, face view. E.—Head of larva, side view. F.—Dorsal view of seventh, eighth, and ninth abdominal segments. G.—Posterior view of segments 7 to 10. 46
10. Pupa and adult of banana root-borer: A.—Ventral view of pupa. B.—Lateral view of head and thorax of pupa. C.—Dorsal view of pupa. 46
11. A.—Young, healthy banana plant bulb with lateral roots. B.—Young banana plant cut into, showing work of the larvæ of the banana root-borer 46

FURTHER STUDIES ON THE INFLUENCE OF HUMIDITY UPON THE STRENGTH AND ELASTICITY OF WOOL FIBER

Text Figures

1. Graphs showing the effect of humidity upon the breaking strength of wool fiber 56
2. Graphs showing the effect of humidity upon the elasticity of wool fiber ... 57
3. Graphs showing the relation between the diameter and the tensile strength of wool fiber 60
4. Graphs showing the effect of humidity upon the breaking strength and elasticity of wool fiber 62

COMPOSITION AND DENSITY OF THE NATIVE VEGETATION IN THE VICINITY OF THE NORTHERN GREAT PLAINS FIELD STATION

Text Figures

1. Diagram of grass mat: A, from side; B, from above. *a*, Basal cover; *b*, foliage cover 68
2. Meter quadrat in 30-acre pasture mapped in detail in 1915 69

Plates

12. General view of native vegetation near Mandan, N. Dak., showing composition and density 72
13. A.—View across area of *Andropogon furcatus*. B.—Close view of *Aristida longiseta* bunches 72
14. A.—Close view, from above, of meter quadrat in 30-acre pasture. B.—Meter quadrat in 30-acre pasture 72

EFFECT OF REACTION OF SOLUTION ON GERMINATION OF SEEDS AND ON GROWTH OF SEEDLINGS

Text Figures

Page

1. A, graph showing the relation of reaction to the contents of $H_3C_6H_5O_7$ and NaOH employed in the cultures of series A; B, graph showing the change of reaction found after 4 days' growth of wheat seedlings in series A... 76
2. A, graph showing the change in reaction obtained by electrometric titration in series B; B, graph showing the change of reaction found after 4 days' growth of wheat seedlings in cultures of series B..... 77
3. Graph showing the relation of growth of wheat seedlings to reaction in series A..... 79
4. Graphs showing the relation of growth of wheat, soybean, corn, and alfalfa seedlings to reaction in series B..... 81
5. Graphs showing the relative weights of sprouts produced by seeds of wheat, corn, soybeans, alfalfa, and red clover in 7-day germination period at various reactions..... 89

Plate

15. A.—Method of growing wheat seedlings. B.—Appearance of wheat seedlings in series A at time of harvesting. C.—Appearance of wheat seedlings in series B at time of harvesting..... 96

PHILIPPINE DOWNY MILDEW OF MAIZE

Text Figures

1. Graph showing variation in length of conidia of *Sclerospora philippinensis*.. 111
2. Graph showing variation in diameter of conidia of *Sclerospora philippinensis*. 111
3. Diagram showing ratios of length to width of conidia of *Sclerospora philippinensis*, arranged in classes and indicating limits of variation and mode.. 111

Plates

A. Young maize plant, showing the effects of a very early attack of the downy mildew on a small, early maturing variety, Manobo Yellow..... 122
B. Young maize plants, showing the effects of later attack of the downy mildew on a large, late-maturing variety, Guam White Dent..... 122
16. A.—Portion of a field of Moro White maize, showing heavy loss from the downy mildew. B.—View near the edge of a field of Guam White Dent maize, showing the ravages of the downy mildew..... 122
17. A.—A frequently encountered type of downy mildew effect. B.—A serious case of downy mildew injury, one of hundreds in a badly attacked maize field..... 122
18. A.—A common result of downy mildew attack. B.—A maize plant seriously injured by the downy mildew stands in front. C.—One hill in a maize plot which lost heavily from attacks by the downy mildew..... 122
19. A.—A case of abnormal growth of a maize plant as the result of an attack by downy mildew. B.—A maize plant which, when nearly mature, became infected by the downy mildew through a small sucker previously developed..... 122
20. A.—A deformed and partly sterile ear complex produced by a maize plant as a result of downy mildew infection. B.—A maize ear developed abnormally as a result of the downy mildew. C.—Ear of a maize plant infected by the downy mildew..... 122

Page

21. A.—A near view of the thick down of conidiophores which has been produced on the upper surface of a badly diseased maize leaf. B.—Upper surface of a badly infected maize leaf from a maturing plant. C.—Upper surface of the middle portion of a maize leaf from a very young plant which has only recently developed the markings of the disease..... 122

22. A.—View of a row of Egyptian sorghum showing tall, green, healthy plants at the left and at the right a dwarfed, yellowish white plant whitch is infected by the downy mildew. B.—Near view of this diseased sorghum plant. C.—A comparative view of healthy teosinte (right), and teosinte seriously infected with the downy mildew (left)..... 122

23. A.—Portion of the typical crooked, irregular mycelum with numerous haustoria which is found in the mesophyll of badly infected leaves, here freed from the host tissue by maceration. B.—Longitudinal section cut from the center of a maize stem 8 inches from the ground. C.—Portion of the mycelium freed by maceration from tissue of the midrib at the base of a badly infected leaf. D.—Hypha cut in cross section as it lies between three adjacent mesophyll cells of the host. E.—Transverse section from a badly infected portion of a maize leaf, showing the abundant mycelium running between the cells of the bundle sheath and forming in the substomatal air chamber the branches (*a*) that grow out through the stoma to form the conidiophores. F.—Portion of a hypha lying between adjacent mesophyll cells, one of which has formed a many-layered wall around the haustorium invading it. G.—Portion of a hypha similar to that shown in F but with the haustorium unhindered in its invasion of the host cell. H.—Bit of mycelium such as is shown in A but more highly magnified to show the haustoria..... 122

24. A.—Slender, sparingly branched conidiophore bearing comparatively few conidia. B.—Tip of branch with two conidia *in situ*. C.—Stout, much-branched, mature conidiophore bearing 38 spores. D.—Upper portion of a nearly mature conidiophore with one secondary branch which has failed to branch further and has terminated in a single conidium only. E.—Small, stunted, sparingly branched conidiophore produced on maize during the light dew of the hot, dry season. F.—Basal cell with two thick crosswalls; from maize. G.—An unusual basal cell with two septa and an abnormally large footlike base. H, J, L.—Typical basal cells of conidiophores. I.—Upper portion of an underdeveloped conidiophore bearing three spores on sterigmata arising directly from the top of the main axis. K.—Tip of an ultimate branch with two sterigmata-bearing conidia. M.—Basal cell of a conidiophore from teosinte with septum formation progressing by the centripetal extension of a cellulose-pectose ring..... 122

25. A.—Conidiophore from sorghum, partly matured and bearing few conidia. B.—Conidiophore from teosinte, nearly mature, with extensive system of branches bearing many conidia. C.—Typical conidia from sorghum. D.—Typical conidia from teosinte. E.—Typical conidia from teosinte which have germinated in dew on the leaf surface. F.—Conidium from teosinte germinating by an extensive branched hypha when maintained in dew at 7° C. G.—Conidium from teosinte germinating while still attached to its sterigma. H.—Typical conidia from maize, showing common variations in shape and size. I.—Two conidia from maize just beginning to germinate in rain water. J.—Two conidia from maize germinating in sterilized brook water maintained at 8° C. K.—Coni-

Page

dium from maize germinating in dew on the leaf surface. L.—Conidium from maize giving rise to extensive branching hyphae in a dilute decoction of young maize kernels 122

EFFECT OF DRUGS ON MILK AND FAT PRODUCTION

Text Figures

1. Graph showing effect of tonic mixture No. 1 on butter-fat and milk yield .. 127
2. Graph showing effect of air-slaked lime on butter-fat and milk yield 127
3. Graph showing effect of Fowler's solution of arsenic on butter-fat and milk yield 128
4. Graph showing effect of powdered gentian on butter-fat and milk yield 128
5. Graph showing effect of the German tonic mixture on butter-fat and milk yield 129
6. Graph showing effect of physostigmine sulphate on butter-fat and milk yield 129
7. Graph showing effect of sodium bicarbonate on butter-fat and milk yield. 130
8. Graph showing effect of ginger on butter-fat and milk yield 130

HALO-BLIGHT OF OATS

Plates

C. Halo lesions on flag leaves of Wisconsin No. 14 oats 172
26. Typical isolated halo lesions 172
27. Halo lesions on Wisconsin No. 14 oats produced by spraying with a water suspension of the stock organism May 26, 1917 172
28. Infection from untreated 1916 seed of Wisconsin No. 124 oats 172
29. Spikelets of Wisconsin No. 14 oats: A.—Left and center spikelets show natural infection with halo-blight. B.—Upper spikelet shows typical isolated halo lesion near base 172
30. A.—Two per cent +5 glucose Difco peptone-beef bouillon agar slant of No. 36. B.—Two per cent potato-dextrose agar slants 172
31. A.—Three-day colony of stock on 2 per cent dextrose-potato agar. B.—Two-day colonies of stock on +10 beef-peptone agar. C.—Five-day colony of stock on potato-dextrose agar. D.—Five-day colony of No. 36 on +10 beef-peptone agar. E.—Three-day colony on 2 per cent glucose Difco peptone beef bouillon agar. F.—Seven-day colony of stock on +15 beef-peptone agar. G.—Margin of 3-day colony of stock on +15 gelatin 172
32. A.—Five-day colonies of stock on potato-dextrose agar. B.—Three-day colony of stock on potato-dextrose agar 172
33. Isolations from sections of halo lesion 172
34. A.—No. 36 from 24-hour potato-dextrose agar slant; carbol fuchsin stain. B. Stock from 24-hour potato-dextrose agar; Ribbert's capsule stain. C.—Stock from 4-day potato-dextrose agar; carbol fuchsin stain, showing long chains. D.—Stock from 3-day potato-dextrose agar; Ribbert's capsule stain. E.—Stock from 1-day +15 beef-peptone agar; Van Ermengem stain. F.—No. 36 from 1-day +5 beef-peptone agar; Caesar-Gil stain .. 172
35. Sections of oat leaves through halo lesions, showing bacteria in the tissues. 172

BEHAVIOR OF THE CITRUS-CANKER ORGANISM IN THE SOIL

Plates

36. A.—Citrus-cankers on mature wood of trunk of *Citrus aurantifolia*. B.—Citrus-cankers on mature wood of branches of *Citrus aurantifolia* 206

Page

37. A.—Results of inoculations with *Pseudomonas citri* upon roots of sweet orange (*Citrus sinensis*). B.—Skeletonized leaves of Ellen grapefruit recovered from buried soil 206

PRODUCTION OF CONIDIA IN GIBBERELLA SAUBINETII

Text Figure

1. Conidial production in *Gibberella saubinetii* (Mont.) Sacc.: A, Ascospores from cornstalk, germinated in distilled water, producing conidia in three days; B, D, typical conidia and conidiophore from a 28-hour-old hanging drop culture from a conidium from A; C, germinating conidia from a 52-hour-old plate culture; E, conidiophore and germinating conidia from a 47-hour-old colony in a Van Tiegham cell 236

EFFECT OF MANURE-SULPHUR COMPOSTS UPON THE AVAILABILITY OF THE POTASSIUM OF GREENSAND

Text Figure

1. Diagrams showing relation of the water-soluble acidity to the water-soluble potassium at different time periods for different greensand composts 250

RUST IN SEED WHEAT AND ITS RELATION TO SEEDLING INFECTION

Text Figure

1. Diagram of air-washing apparatus for isolated room used for growing rust-infected seed 268

Plates

38. Heads of Kubanka durum wheat heavily infected with stemrust 277
39. A.—Portion of one of the heads shown in Plate 38. B.—Wheat kernels showing typical stemrust infection 278
40. Exterior view of isolated room in the pathological greenhouse at the University of Wisconsin, showing (*a*) the exterior portion of air-washing apparatus used to wash all air drawn into the room, (*b*) the canvas curtain used for shading on warm days, and (*c*) the sprinkling attachment used to throw spray of water over the roof to aid in keeping the room cool 278
41. A.—Photograph of wheat grown in flats in isolated room in greenhouse at the University of Wisconsin. B.—Same plants as A, when well headed 278
42. Longitudinal section through hilar portion of an immature wheat kernel, showing sorus of stemrust 278
43. Longitudinal section through the hilum of a wheat kernel infected with stemrust, showing unusually large internal sori extending nearly across the kernel 278
44. Cross section of a mature wheat kernel infected with stemrust, showing telia in the ventral groove 278
45. Enlarged portion of section shown in Plate 44, showing telia on surface of ventral groove 278
46. Longitudinal section of embryo of germinated wheat kernel showing large internal rust at *x* in hilar tissue at base of embryonic tissue 278
47. Longitudinal section of the embryo further advanced in development than that shown in Plate 50 278
48. Longitudinal section through young secondary root of wheat embryo, showing presence of intracellular mycelium 278

PRACTICAL UNIVERSALITY OF FIELD HETEROGENEITY AS A FACTOR INFLUENCING PLOT YIELDS

Text Figures

Page

1. Montgomery's diagram of 5.5 by 5.5 foot plots of Turkey wheat, showing variations in the percentage of nitrogen in the grain 283
2. Diagram showing yield of alfalfa in first cutting, 1913, on the Huntley experimental tract 286
3. Diagram showing yield of alfalfa in second cutting, 1913, on the Huntley experimental tract 287
4. Diagram showing yield of alfalfa in first cutting, 1914, on the Huntley experimental tract 288
5. Diagram showing yield of alfalfa in second cutting, 1914, on the Huntley experimental tract 289
6. Diagram showing yield of alfalfa in third cutting, 1914, on the Huntley experimental tract 290
7. Diagram showing yield of unhusked rice in Coombs and Grantham's 54 plots ¼ by ¼ chain square 296
8. Diagram showing yield of ear corn, 1915, on the Huntley experimental tract. 297
9. Diagram showing yield of ear corn, 1916, on the Huntley experimental tract. 298
10. Diagram showing location of sample areas examined for soil moisture in a field at San Antonio Experimental Farm 302

TRANSMISSION OF THE MOSAIC DISEASE OF IRISH POTATOES

Plates

49. Vines of Green Mountain variety inoculated with juice from healthy foliage of the same variety 338
50. Vines of Bliss Triumph variety inoculated with juice from healthy foliage of Irish Cobbler variety 338
51. Vines of Irish Cobbler variety inoculated with juice from healthy foliage of the same variety 338
52. Vines of Green Mountain variety inoculated with juice from mosaic foliage of the same variety 338
53. Vines of Green Mountain variety inoculated with juice from mosaic foliage of Bliss Triumph variety 338
54. Vines of Green Mountain variety inoculated with juice from mosaic foliage of Irish Cobbler variety 338
55. Vines of Bliss Triumph variety inoculated with juice from mosaic foliage of Irish Cobbler variety 338
56. Vines of Irish Cobbler variety inoculated with juice from mosaic foliage of the same variety showing bad stage of mosaic 338

RELATIVE SUSCEPTIBILITY TO CITRUS-CANKER OF DIFFERENT SPECIES AND HYBRIDS OF THE GENUS CITRUS, INCLUDING THE WILD RELATIVES

Plates

57. Leaf of *Casimiroa edulis* with naturally occurring spots from the greenhouse inoculations 362
58. Infected leaves from plants of *Chaetospermum glutinosum* in the greenhouse experiments, showing the types of canker spots produced 362

Page

59. A.—Leaves of *Atalantia citrioides* and *A. ceylonica* (center) from plants in the greenhouse experiments, showing the canker spots typically produced on these plants. B.—Compound leaf of *Hesperthusa crenulata* from isolation field, with naturally occurring canker spots on two of the leaves. C.—Leaves of *Microcitrus Garrowayi* from plants in the greenhouse experiments, with different types of canker spots. D, E.—Infected leaves and twigs of *Eremocitrus glauca* from greenhouse plants, showing the large flat spots on the leaves and the rather corky spots on the twig and thorn 362
60. A.—Typically infected leaf of *Fortunella margarita*. B.—Old leaf of *Citrus grandis*, with raised, compact, oily, unruptured spots. C.—*Fortunella Hindsii*, with ruptured corky spots. D.—*Citrus* sp., Kansu of Yuzu Orange. E.—*Citrus aurantifolia*, showing typical infection 362
61. Upper and lower leaf surfaces of a *Citrus hystrix* leaf with a heavy natural canker infection 362
62. Typically infected leaves of *Citrus grandis* from field plants showing extreme susceptibility 362
63. A.—Leaves and twigs of faustrime from greenhouse experiment with typical spots. B.—Types of spots found on *Citrus nobilis* (King of Siam, Naranjita, and tangerine). C.—Leaf of the citrangequat from greenhouse experiment. D.—Citrumelo leaf with typical canker spots. E.—Upper and lower surface of a naturally infected leaf of *Citrus mitis* in the field .. 362
64. Naturally infected leaves of *Citrus nobilis* var. *unshiu* from the field, showing various types of spots produced 362
65. Some of the hybrids of *Porcirus trifoliata*, showing vigor, type of growth, leaf characters, and relative susceptibility to citrus-canker, arranged in order of their susceptibility 362
66. A.—Limelos in the greenhouse inoculation experiments, showing type of growth, leaf characters, and susceptibility to citrus-canker. B.—Limequats in the greenhouse inoculation experiments, showing type of growth, character of leaves, and susceptibility to citrus-canker 362
67. A.—Siamelos in the greenhouse inoculation experiments, showing type of growth, leaf characters, and susceptibility to citrus-canker. B.—Comparison of type of growth, leaf characters, and susceptibility to citrus-canker in clemelo, satsumelo, and tangelo in the greenhouse inoculation experiments 362
68. A.—Results of the greenhouse inoculations with some of the false hybrids. B.—Tangelos in the greenhouse inoculation experiments, showing type of growth, leaf character, and susceptibility to citrus-canker 362

PRESOAK METHOD OF SEED TREATMENT: A MEANS OF PREVENTING SEED INJURY DUE TO CHEMICAL DISINFECTANTS AND OF INCREASING GERMICIDAL EFFICIENCY

Text Figures

1. Graph showing effect of formalin 1 to 400 treatments with and without presoaking 370
2. Graph showing effect of formalin 1 to 200 treatments with and without presoaking 372
3. Graph showing effect of formalin 1 to 320, with and without presoaking 374
4. Graph showing effect of copper sulphate 1 to 80, with and without presoaking, on wheat and barley seed germination 375
5. Graph showing effect of formalin 1 to 320, with and without presoaking, on wheat seed germination under field conditions 377

Page

6. Graph showing effect of formalin and copper-sulphate presoak treatments of ½-bushel wheat seed lots 378
7. Graph showing effect of formalin 1 to 320 treatments for various periods, with and without presoaking 385
8. Graph showing effect of formalin 1 to 320 and 1 to 200 on germination of corn, barley, and oats with and without presoaking 386
9. Curve showing rate of absorption of water by dry wheat seeds 387

Plates

69. Relative injury to wheat-seed germination caused by short and long formalin treatments 392
70. Effect of formalin 1 to 400 treatment for 6 hours, with and without 6-hour presoak. 1. Fultz wheat. 2. Poole wheat 392
71. Effect of formalin 1 to 400 treatment for 6 hours, with and without 6-hour presoak. 1. Fulcaster wheat. 2. Turkey wheat 392
72. Stimulating effect of the presoak method of treatment with formalin 1 to 400 392
73. Effect of formalin 1 to 200 treatment for 6 hours, with and without 6-hour presoak 392
74. Effect of formalin 1 to 320 treatment for 6 hours, with and without 6-hour presoak 392
75. 1. Fife wheat: A, B, control, 83 per cent germination; C, D, seeds treated with formalin 1 to 320 for 3 hours, 62 per cent germination; E, F, seeds presoaked 6 hours, then formalin 1 to 320 for 6 hours, 90 per cent germination. 2. Poole wheat: A, B, control, 85 per cent germination; C. D. seeds treated with formalin 1 to 320 for 6 hours, 55 per cent germination; E, F, seeds presoaked 6 hours, then formalin 1 to 320 for 6 hours, 88 per cent germination. 3. Effect of soaking in water throughout presoak period, compared with procedure of keeping moist 6 hours: A, B, Fife wheat; C, D, Poole wheat; A, C, seeds soaked in water 5 hours, then treated with formalin 1 to 320 for 7 hours; B, D, seeds soaked in water 10 minutes, drained, and kept moist 6 hours, then treated with formalin 1 to 320 for 6 hours 392
76. Effect of formalin and copper sulphate on wheat and of copper sulphate on barley, with and without presoaking 392
77. Effect of presoak method used with copper-sulphate treatment of wheat 392
78. Effect of presoak method used with formalin 1 to 320 on wheat under field conditions 392
79. Effect of presoak method used with formalin 1 to 400 on blackchaff bacteria 392
80. Effect of presoak method used with formalin 1 to 400 on blackchaff bacteria 392
81. Effect of presoak method on barley and oats 392
82. Effect of presoak method used on barley, oats, and corn 392

DAILY DEVELOPMENT OF KERNELS OF HANNCHEN BARLEY FROM FLOWERING TO MATURITY AT ABERDEEN, IDAHO

Text Figures

1. Graph showing length, lateral diameter, and dorsoventral diameter of barley kernels for the 25 days following flowering 395
2. Graph showing percentage of moisture per kernel from date of flowering 397
3. Graph showing maximum and mean daily temperatures recorded at Aberdeen, Idaho, from July 8 to August 4, 1916 (broken line), and from July 15 to August 10, 1918 (solid line) 398

Page

4. Graph showing lateral and dorsoventral diameters of the ovary tip as compared with length, lateral diameter, and dorsoventral diameter of the kernel for the 26 days following flowering in 1916 408
5. Graph showing average length of kernels 6, 7, and 8 (solid line), average lateral diameter (dotted line), and average dorsoventral diameter (broken line) from plot 1 in 1917 410
6. Graph showing average length of barley kernels, including ovary tip, from flowering to near maximum development in plot 1 in 1917 411
7. Graph showing average lateral diameter of barley kernels from flowering until near maximum development, plot 1, 1917 412
8. Graph showing dorsoventral diameters of barley kernels from flowering until near maximum development 413
9. Graph showing wet weight of individual kernels 5, 8, and 10, by days, from date of flowering to near maturity in 1917 418
10. Graph showing average wet weights of kernels from flowering to maturity in plot 1 in 1917 419
11. Graph showing dry matter per kernel from date of flowering to near maturity in 1916 (dotted line) and in 1917 (solid line) 420
12. Graph showing dry matter per kernel at 12-hour intervals from flowering to maturity 421
13. Graph showing dry-weight gain of kernels 6, 7, and 8 in 12-hour periods .. 422
14. Graph showing percentage of moisture in morning and evening samples of Hannchen barley in 1917 422
15. Graph showing nitrogen per kernel from date of flowering in 1916 (broken line) and in 1917 (solid line) 424
16. Graph showing ash per kernel from date of flowering in 1916 (broken line) and in 1917 (solid line) and the percentage of ash in 1917 (dotted line) .. 426
17. Periods of development of the barley kernel as indicated by records during three years at Aberdeen, Idaho 427

Plates

83. A.—Fertilized ovary. B.—Kernel 1 day old. C.—Kernel 2 days old. D.—Kernel 3 days old 430
84. A.—Kernel 4 days old. B.—Kernel 5 days old. C.—Kernel 6 days old. D.—Kernel at later stage of development 430
85. Kernel 5 days after fertilization 430
86. Kernel 6 days after fertilization 430
87. Kernel 9 days after fertilization 430
88. Kernel 14 days after flowering 430
89. Kernel 20 days after fertilization, at which time growth was nearly completed 430
90. Kernel 25 days after fertilization, growth completed 430
91. Section of a nearly mature kernel, showing cells next the furrow 430

DEVELOPMENT OF BARLEY KERNELS IN NORMAL AND CLIPPED SPIKES AND THE LIMITATIONS OF AWNLESS AND HOODED VARIETIES

Text Figures

1. Graph showing growth in length, lateral diameter, and dorsoventral diameter of kernels of Manchuria barley in normal and clipped spikes 448
2. Graph showing wet weight of kernels of Manchuria barley from normal and clipped spikes 449

Page

3. Graph showing dry matter in kernels of Manchuria barley from normal and clipped spikes 449
4. Graph showing total ash in kernels of Manchuria barley from normal and clipped spikes 450
5. Graph showing total nitrogen in kernels of Manchuria barley from normal and clipped spikes 451
6. Graph showing water in kernels of Manchuria barley from normal and clipped spikes 451
7. Graph showing growth in length, lateral diameter, and dorsoventral diameter of kernels of Hannchen barley in normal and clipped spikes 463
8. Graph showing wet weight of kernels of Hannchen barley from normal and clipped spikes 464
9. Graph showing dry matter in kernels of Hannchen barley from normal and clipped spikes 465
10. Graph showing percentage of ash in the kernels, rachises, paleae, and awns of normal spikes of Hannchen barley and in the kernels, rachises, and paleae of clipped spikes 467
11. Graph showing total nitrogen in kernels of Hannchen barley from normal and clipped spikes 468
12. Graph showing water in kernels of Hannchen barley from normal and clipped spikes 469
13. Graph showing relation of length of awn to weight of clipped kernels and unclipped spikelets on a 2-rowed barley grown at Arlington Farm, Va. . 470

INVESTIGATIONS IN THE RIPENING AND STORAGE OF BARTLETT PEARS

Text Figures

1. Sugars in Bartlett pears from Sacramento, Calif 484
2. Sugars in Bartlett pears from Suisun, Calif 485
3. Sugars in Bartlett pears from Medford, Oreg 486
4. Sugars in Bartlett pears from Yakima, Wash 487
5. Acids in Bartlett pears from Sacramento, Calif 490
6. Acids in Bartlett pears from Suisun, Calif 491
7. Acids in Bartlett pears from Medford, Oreg 491
8. Acids in Bartlett pears from Yakima, Wash 492

FURTHER DATA ON THE ORANGE-RUSTS OF RUBUS

Plates

D. 1.—Infected black raspberry leaf covered with the caeomas of *Gymnoconia interstitialis*. 2.—Blackberry leaf infected with the short-cycled orange-rust 512
92. Manner of germination of the spores of the two orange-rusts 512
93. Short-cycled orange-rust 512
94. *Gymnoconia interstitialis* 512

MOSAIC DISEASE OF CORN

Plates

95. Mosaic disease of corn 522
96. Mosaic disease of corn: Effect of early infection on the ear 522

GENETICS OF RUST RESISTANCE IN CROSSES OF VARIETIES OF TRITICUM VULGARE WITH VARIETIES OF T. DURUM AND T. DICOCCUM

Plates

Page

97. A.—Pollen grains of Marquis wheat B. —F_1 Marquis×Kubanka (C I 2094), showing sterile grains C.—Pollen grains of Kubanka (C I 2094). D.—Stems of Kubanka (C I 2094) grown under rust-epidemic conditions. E.—F_1 Kubanka (C I 2094)×Marquis, showing normal uredinia. F.—Marquis, the susceptible parent. H.—F_1 emmer (Minnesota 1165)×Marquis, showing no normal uredinia. I.—Minnesota 1165, the resistant emmer parent 542

98. A, B, C.—Face and side views, respectively, of heads of Iumillo (C I 1736), F_1 Iumillo×Marquis, and Marquis. D, E, F.—Face and side views, respectively, of heads of emmer, Minnesota 1165, F_1 emmer×Marquis, and Marquis. G, H, I.—Kernels of Marquis, F_1 emmer×Marquis, and emmer 542

99. Representative heads of F_3 families of the cross between durum and Marquis. 542

100. A.—F_3 family of a cross between emmer (Minnesota 1165), and Marquis, showing face and side view. B.—Heads of an F_3 family which resembled common wheat. C.—Heads of different plants of an F_3 family of a cross between emmer (C I 1524) and Marquis. D.—Represents a very frequent sort of segregation obtained in the F_3 generation 542

101. Heads of resistant and susceptible wheat obtained in the F_3 generation from the cross between Marquis and durum 542

102. Greenhouse experiment 542

OCCURRENCE OF THE FIXED INTERMEDIATE, HORDEUM INTERMEDIUM HAXTONI, IN CROSSES BETWEEN H. VULGARE PALLIDUM AND H. DISTICHON PALMELLA

Plates

103. Individual heads, representing the three phenotypic progeny classes in which the lateral florets bear awns 591

104. Individual heads, representing the phenotypic progeny classes in which the lemmas of the lateral florets are rounded and awnless 591

105. A.—Awn-pointed individual, heterozygous for regressive 6-rowed and 2-rowed characters. B.—Short-awned individual, heterozygous for regressive 6-rowed and 2-rowed characters. C.—Long-awned individual, heterozygous for regressive 6-rowed and 2-rowed characters. D.—Three spikes of Mansfield barley from the same plant, showing the variations of fertility in the lateral florets 591

106. A.—Infertile spike of potentially fertile *Hordeum intermedium*. B.—Fertile spike of *H. intermedium*. C.—Var. *atterbergii*, probably a sterile *intermedium* 591